



ALTAIR Technical Note X version 1

Calculations of the Output of a Diffused Planar Light Source

Justin Albert
jalbert@uvic.ca

Jan. 2015

This report provides analytic calculations of angular and position-space distributions of the output flux from a diffused planar light source, one example being a light source, such as a laser or LED, which is shone through a flat sheet of translucent plastic. Both simple small-angle scattering calculations, and more complex large-angle scattering formulae, are provided. Comparisons of these calculated analytic formulae with results from simple numerical Monte Carlo simulations are additionally provided.

Contents

1	Initial Assumptions and Examples	1
1.1	The “succession-of-very-thin-layers” (SOVTL) approximation .	1
1.2	The “scattering centres” (SC) approximation	3
2	Calculations of Small-Angle Scattering Formulae	5
2.1	Three spatial dimensions	5
2.2	Deviation from normal incidence angle	5
2.3	Calculation using non-projected “ δy ” rather than dy	5
2.4	Adding absorption	5
2.5	Impact-parameter-dependent SC approximation scattering . .	5
2.6	Initial considerations for non-Gaussian scattering: the exam- ple of Rayleigh scattering	5
3	Calculations of Large-Angle Scattering Formulae	6
3.1	Non-Gaussian scattering redux	6
3.2	Usage of Bayes’ Theorem	6
3.3	Large-angle scattering with absorption	6
4	A Toy Monte Carlo Simulation and Its Results	7
4.1	Description of the Toy MC	7
4.2	Results with Gaussian scattering	7
4.3	Results with Rayleigh scattering	7
5	Comparison of Analytic Calculation Formulae with Toy Monte Carlo Results	8
A	Some of the Necessary Input Formulae	9
A.1	A proof of the Bienaymé formula	9
A.2	The moments and bivariate moments of Gaussian distributions	9
B	A Brief List of the Important Results Found	10
	References	11

1 Initial Assumptions and Examples

When performing analytical, and even numerical, calculations of photon scattering and absorption in a medium, clearly a number of assumptions must be made in order to develop a model and perform such calculations. In addition to typical approximations we shall use at various points (but not universally, we will carefully note each approximation we use in any given calculation) — two examples being the small angle approximation, and an assumption that scattering angles are Gaussian — we will extensively utilize two approximate models for the optical behaviour of a sheet of diffusive material, which will be denoted as the “succession-of-very-thin-layers” approximation and the “scattering centres” approximation. These two approximate models are described below, each with an example calculation utilizing the given model.

1.1 The “succession-of-very-thin-layers” (SOVTL) approximation

We first consider diffusive photon scattering in the SOVTL approximation as shown in Fig. 1.1 at the top of the following page.

Consider non-absorptive scattering in this approximation, where each of the scattering angles, as described in the figure caption as having RMS equal to $\phi_{\text{RMS}}\sqrt{\frac{dx}{t_0}}$, is additionally small (*i.e.*: $\phi_{\text{RMS}}\sqrt{\frac{dx}{t_0}} \ll \pi/2$) and is Gaussian-distributed with mean 0. Let us calculate 1) the RMS size of the output beam, *i.e.* the distribution of output positions of photons exiting the upper surface of the scattering medium, and also 2) the distribution of output angles of these same exiting photons.

The projected scattering distance dy from a given scatter from a layer at position x must equal $(s-x)[\tan(\Phi_{\text{sc}}+\phi_{\text{sc}})-\tan(\Phi_{\text{sc}})]$, which is approximately equal to $(s-x)[\Phi_{\text{sc}}+\phi_{\text{sc}}-\Phi_{\text{sc}}] = (s-x)\phi_{\text{sc}}$, since we are assuming all ϕ_{sc} , and thus Φ_{sc} , are small. Thus, if we consider the distribution of possible scattering distances dy due to the scatter at x , the variance of that distribution $\text{var}(dy)$ must equal $(s-x)^2\text{var}(\phi_{\text{sc}})$, since we are treating s and x as constants here. We know (per the SOVTL approximation) that $\text{var}(\phi_{\text{sc}}) = \phi_{\text{RMS}}^2 \frac{dx}{t_0}$, and thus $\text{var}(dy) = (s-x)^2\text{var}(\phi_{\text{sc}}) = (s-x)^2\phi_{\text{RMS}}^2(\frac{dx}{t_0})$. We would like to calculate 1)




Figure 1.1: The “succession-of-very-thin-layers” (SOVTL) approximation to photon scattering in a diffusive medium. Photons in the beam can scatter at the location of each of the thin layers in the medium. We let the RMS angle of scattering for each very thin layer of thickness dx — with an example scatter being given by ϕ_{sc} above — be $\phi_{\text{RMS}}\sqrt{\frac{dx}{t_0}}$, where ϕ_{RMS} and t_0 are material-dependent parameters. The (signed) sum of all scattering angles prior to ϕ_{sc} is denoted as Φ_{sc} . The distance dy is the projected scattering distance on the upper surface of the medium due only to the single scatter ϕ_{sc} . If we are including absorption effects, we may also let the probability of non-absorption in a given layer be $e^{-dx/\tau}$, where τ is another material-dependent parameter.

$\text{var}(y)$, where $y = \sum_{\text{layers}} dy = \int_{x=0}^{x=s} dy$; as the square root of $\text{var}(y)$ will be the RMS size of the output beam, and also 2) $\text{var}(\sum_{\text{layers}} \phi_{sc})$. We have that $\text{var}(y) = \text{var}[\sum dy] = \text{var}[\int dy]$. From the “Bienaymé formula” (see Appendix A for a proof), we have that $\text{var}[\sum_i x_i] = \sum_i \text{var}(x_i)$ for any set of uncorrelated variables x_i , thus $\text{var}(y) = \text{var}[\int dy] = \int \text{var}(dy) = \int_0^s (s-x)^2 \phi_{\text{RMS}}^2 (\frac{dx}{t_0}) = \frac{\phi_{\text{RMS}}^2}{t_0} \int_0^s (s-x)^2 dx = \frac{\phi_{\text{RMS}}^2}{t_0} \left[-\frac{1}{3}(s-x)^3 \right]_0^s = \frac{\phi_{\text{RMS}}^2 s^3}{3t_0}$. Thus, 1) the RMS size of the output beam $= \sqrt{\text{var}(y)}$ must be the square root of this, *i.e.* $\frac{\phi_{\text{RMS}} s^{3/2}}{\sqrt{3t_0}}$.

To find 2), we use Bienaymé again and thus note that $\text{var}(\sum_{\text{layers}} \phi_{sc}) =$

$\sum_{\text{layers}} \text{var}(\phi_{\text{sc}}) = \int_{x=0}^{x=s} \text{var}(\phi_{\text{sc}})$. This is given by $\int_{x=0}^{x=s} \phi_{\text{RMS}}^2(\frac{dx}{t_0}) = \frac{\phi_{\text{RMS}}^2}{t_0} \int_{x=0}^{x=s} dx = \frac{\phi_{\text{RMS}}^2 s}{t_0}$. Thus the standard deviation of the output angles is the square root of this, *i.e.* $\phi_{\text{RMS}} \sqrt{\frac{s}{t_0}}$.

Formulae for angular and spatial output distributions for more complex cases than the above in the SOVTL approximation will be calculated in Chapters 2 and 3.

1.2 The “scattering centres” (SC) approximation

We next consider diffusive photon scattering in the SC approximation as shown in Fig. 1.2 below:




Figure 1.2: The scattering centres (SC) approximation to photon scattering. Photons in the beam can scatter at the random location of each of the scattering centres in the medium. We let the mean number of scattering centres per unit area be n (with $ns^2 \gg 1$). Let the RMS angle of scattering for each centre be ϕ_{RMS} , and let the (circular) scattering cross-section of each centre be t_0 . We initially consider scattering to be impact parameter independent within the circular cross-section, and to have a Gaussian scattering angle distribution.

Consider non-absorptive scattering as in the previous section, where each of the scattering angles is small (*i.e.*: $\phi_{\text{RMS}} \ll \pi/2$) and is Gaussian-distributed with mean 0. We calculate (as above in the SOVTL approximation) 1) the RMS size of the output beam, and 2) the distribution of output angles of the same exiting photons.

The projected scattering distance dy from a given scatter from a centre at position x is (as with SOVTL) approximately equal to $(s-x)\phi_{\text{sc}}$. Thus, if we consider the distribution of possible scattering distances dy due to the scatter at x , the variance of that distribution $\text{var}(dy)$ must equal $(s-x)^2 \text{var}(\phi_{\text{sc}}) = (s-x)^2 \phi_{\text{RMS}}^2$. Using Bienaymé here as well, we have that $\text{var}(y) = \text{var}[\sum dy] = \sum \text{var}(dy) = \sum (s-x)^2 \phi_{\text{RMS}}^2 = \int_0^s (s-x)^2 \phi_{\text{RMS}}^2 \delta i dx$, where δi represents the number of centres hit (or the “chance of hitting a centre”) per the short distance of travel dx . This quantity δi will equal $\frac{(\text{length of travel}) \times t_0 \times n}{dx}$. The length of travel will equal $\frac{dx}{\cos \Phi_{\text{sc}}}$, where Φ_{sc} is the sum of all scattering angles ϕ_{sc} for all scatters prior to the scatter under consideration. Thus we have that $\delta i = \frac{t_0 \times n}{\cos \sum \phi_{\text{sc}}}$. Using the small angle approximation, $\cos \sum \phi_{\text{sc}} \approx 1$, thus $\text{var}(y) \approx \int_0^s (s-x)^2 \phi_{\text{RMS}}^2 t_0 n dx = \phi_{\text{RMS}}^2 t_0 n \left[-\frac{1}{3}(s-x)^3 \right]_0^s = \frac{\phi_{\text{RMS}}^2 t_0 n s^3}{3}$. Thus,

1) the RMS size of the output beam = $\phi_{\text{RMS}} \sqrt{\frac{t_0 n s^3}{3}}$.

To find 2), we use Bienaymé again and thus note that $\text{var}(\sum \phi_{\text{sc}}) = \sum \text{var}(\phi_{\text{sc}}) = \int_{x=0}^{x=s} \text{var}(\phi_{\text{sc}}) = \int_{x=0}^{x=s} \phi_{\text{RMS}}^2 \delta i dx = \phi_{\text{RMS}}^2 t_0 n \int_{x=0}^{x=s} dx = \phi_{\text{RMS}}^2 t_0 n s$. Thus the standard deviation of the output angles is the square root of this, *i.e.* $\phi_{\text{RMS}} \sqrt{t_0 n s}$.

2 Calculations of Small-Angle Scattering Formulae

- 2.1 Three spatial dimensions
- 2.2 Deviation from normal incidence angle
- 2.3 Calculation using non-projected “ δy ” rather than dy
- 2.4 Adding absorption
- 2.5 Impact-parameter-dependent SC approximation scattering
- 2.6 Initial considerations for non-Gaussian scattering: the example of Rayleigh scattering

3 Calculations of Large-Angle Scattering Formulae

3.1 Non-Gaussian scattering redux

3.2 Usage of Bayes' Theorem

3.3 Large-angle scattering with absorption

4 A Toy Monte Carlo Simulation and Its Results

4.1 Description of the Toy MC

4.2 Results with Gaussian scattering

4.3 Results with Rayleigh scattering

5 Comparison of Analytic Calculation Formulae with Toy Monte Carlo Re- sults

A Some of the Necessary Input Formulae

A.1 A proof of the Bienaymé formula

A.2 The moments and bivariate moments of Gaussian distributions

B A Brief List of the Important Results Found

Bibliography

- [1] T. D. Lee and C. N. Yang, Phys. Rev. **104** (1956) 254.
- [2] C. S. Wu *et al.*, Phys. Rev. **105** (1957) 1413.
- [3] J.H. Christensen, J.W. Cronin, V.L. Fitch, and R. Turlay, Phys. Rev. Lett. **13** (1964) 138.
- [4] M. Kobayashi and T. Maskawa, Prog. Th. Phys. **49** (1973) 652.
- [5] V. Fitch, in Nobel Lectures (Physics) World Scientific, Singapore (1991), lecture for 1980 prize.
- [6] I. I. Bigi *et al.*, in *CP Violation*, ed. C. Jarlskog, World Scientific, Singapore (1992).
- [7] M. Gell-Mann and A. Pais, Phys. Rev. **97** (1955) 1387.
- [8] M. Peskin and D. Schroeder, An Introduction to Quantum Field Theory Addison-Wesley, New York (1995).
- [9] R. Aleksan, Quark Mixing and CP Violation, Proceedings of the NATO-ASI Virgin Islands Summer School (1995).
- [10] S. L. Glashow, J. Iliopoulos, and L. Maiani, Phys. Rev. D **2** (1970) 1285.
- [11] A. D. Sakharov, *ZhETF Pis. Red.* **5**, 32 (1967); *JETP Lett.* **5**, 24 (1967).
- [12] V. A. Kuzmin, V. A. Rubakov and M. Shaposhnikov, *Phys. Lett. D* **155** (1985) 36.
- [13] S. Adler, Phys. Rev. **117** (1969) 2426.
- [14] J. Bell and R. Jackiw, Nuo. Cim. **A 60** (1969) 47.
- [15] G. 't Hooft, Phys. Rev. D **14** (1976) 3432.
- [16] M.B. Gavela *et al.*, Nucl. Phys. B **430** (1994) 345.

- [17] The *BABAR* Collaboration, Nucl. Instr. and Methods A **479** (2002) 1.
- [18] B. Aubert *et al.* (The *BABAR* Collaboration), hep-ex/0203008, accepted for publication in Phys. Rev. Lett. .
- [19] Y. Grossman and M. Worah, Phys. Lett. B **395**, 241 (1997),
R. Fleischer, Int. Jour. Mod. Phys. **A 12**, 2459 (1997)
- [20] E. Lipeles, *et al.*, Phys. Rev. D **62** (2000).
- [21] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
- [22] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. **B185**, 218 (1987).
- [23] P. H. Harrison and H. R. Quinn, eds. “The *BABAR* Physics Book”, SLAC-R-504 (1998).
- [24] Particle Data Group. D. E. Groom *et al.*, “Review of Particle Physics”, Eur. Phys. Jour. C **15**, 1 (2000).
- [25] C. Hearty, “Measurement of the Number of $\Upsilon(4S)$ Mesons Produced in Run 1 (B Counting)”, *BABAR* Analysis Document 134, (2001).
- [26] BReco AWG, “Exclusive Reconstruction of Hadronic B Decays to Open Charm”, *BABAR* Analysis Document 150, (2001).
- [27] <http://www.slac.stanford.edu/BFR00T/www/Physics/BaBarData/GoodRuns/dataSets.html>
- [28] <http://www.slac.stanford.edu/BFR00T/www/Physics/Tools/BetaTools/MicroKilling.html>
- [29] private communication from Prof. Aaron Roodman
- [30] E. Varnes “Measurement of the Tracking Efficiency Using 3+1 tau Events”, *BABAR* Analysis Document 87, (2000).
<http://babar-hn.slac.stanford.edu:5090/HyperNews/get/physAnal/1037.html>
<http://babar-hn.slac.stanford.edu:5090/HyperNews/get/physAnal/1024.html>
<http://www.slac.stanford.edu/BFR00T/www/Physics/TrackEfficTaskForce/Recipe/TrackingEfficiencies.html>
- [31] <http://babar-hn.slac.stanford.edu:5090/HyperNews/get/physAnal/1042.html>
- [32] <http://babar-hn.slac.stanford.edu:5090/HyperNews/get/physAnal/1028.html>

- [33] D. Kirkby *et al.* “A User’s Guide to the RooFitTools Fitting Package”, *BABAR* Analysis Document 12 (2000).
- [34] N. Gehrels, *ApJ*, **303**, 336 (1986).
- [35] K. Abe *et al.*, Belle-Conf-0104 (2001).
- [36] PDG 2000.
- [37] B.A.D. 205 Supporting document for summer 2001 sin2beta analysis.
- [38] B.A.D. 356 Supporting document for the Moriond, 2002 sin2beta analysis.
- [39] B. Aubert *et al.* (The *BABAR* Collaboration), hep-ex/0201020, accepted for publication in *Phys. Rev. D*.
- [40] R. Aleksan *et al.*, *Nucl. Phys. B* **361** (1991) 141.
- [41] R. Aleksan *et al.*, *Phys. Lett. B* **317** (1993) 173.
- [42] <http://babar-hn.slac.stanford.edu:5090/HyperNews/get/sin2beta/198.html> (R. N. Cahn).
- [43] J. D. Richman, Heavy-Quark Physics and *CP* violation (Les Houches lectures) (1998).
- [44] B.A.D. 125 Measurement of B^0 mixing using fully reconstructed hadronic B^0 decays
- [45] Defined as $A \sim \sqrt{1 - (m_{ES}/m_0)^2} \times \exp(\kappa(1 - (m_{ES}/m_0)^2))$ for $m_{ES} < m_0$. The default value of m_0 is 5.291 GeV.
- [46] B.A.D. 56 (contact D. Wright) Analysis of $B^0 \rightarrow J/\psi K_L^0$
- [47] B.A.D. 119 (contact J. Beringer) B Tagging in BaBar: Status for the sin2beta Journal Publication
- [48] B.A.D. 102 (contact F.Martinez Vidal) The BaBar Vertexing
- [49] B.A.D. 130 (contact F.Martinez Vidal) Performances and control samples of the *BABAR* vertexing

- [50] B.A.D. 99 (contact O. Long) Measurement of $\sin 2\beta$ with the decay mode $B^0 \rightarrow J/\psi K_L^0$
- [51] B.A.D. 151 (S. Plaszczynski, L. Roos, M.H. Schune) Measurement of the mistag fractions with counting methods
- [52] J. Beringer, *Changes in tagging and deltaT for events used for Osaka results*, sin2beta HN 192, January 19, 2001
- [53] J. Beringer, *Comparison with and Unblinding of the Osaka Subsample*, sin2beta meeting, January 22, 2001
- [54] J. Beringer et al., *Cut Based Tagging*, BaBar Analysis Document 118, Version 1, (2000)
- [55] F. Porter, Interval Estimation Using the Likelihood Function.
- [56] <http://babar-hn.slac.stanford.edu:5090/HyperNews/get/physAnal/1023.html> (A. Gritsan).
- [57] B.A.D. 153 (contact H. Lacker) Constraining the CKM Matrix
- [58] B.A.D. 92 (contact G. Eigen) Inputs for the Unitarity Triangle Fits
- [59] BaBar CKM Working Group,
<http://babar-hn.slac.stanford.edu:5090/HyperNews/get/ckmfit/5.html>
- [60] D.N. Brown, E. Charles, and D. Roberts, “The BaBar Track Fitting Algorithm”, submitted to the CHEP 2000 conference (Berkeley, Jan. 2000)
- [61] F. Frühwirth, “Applications of Filter Method to the Reconstruction of Tracks and Vertices in Events of Experimental High Energy Physics”, HEPHY-PUB 516/88 (Vienna, Dec. 1988)
- [62] R. Cousins *et al.*, “Kalman Filter Track Fits and Track Breakpoint Analysis”, submitted to Nucl. Instr. Meth. (Geneva, Dec. 1999)
- [63] G. Lynch *et al.*, “BaBar Drift Chamber Tracking Conventions”, BaBar Note 488, TNDC Note 99-95 (Stanford, Apr. 1999)